



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$P_2 = \frac{1}{\pi^2} \int_0^{2\sin^{-1}\frac{1}{2}} (\pi - \phi_1) d\theta.$$

This is not integrable in general terms but its value may be readily approximated by methods of mechanical quadrature.

II. Solution by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics in Russell College, Lebanon, Va., and J. A. MOORE, Ph. D., Professor of Mathematics, Millsaps College, Jackson, Miss.

Let $AO = a$, then $CO = \frac{3}{2}a$.

(1). Let his first step place him at the point A , then in order that he may step outside on the second step he must step somewhere on the arc CDB .

Let $AE = t$, $EC = u$, $\angle DAC = \beta$, $P =$ chance in (1), $p =$ chance in (2).

Now $u^2 = a^2 - t^2 = \frac{9}{4}a^2 - (a+t)^2$, $\therefore t = \frac{1}{4}a$.

$\therefore \cos \beta = \frac{1}{4}$.

$\therefore P = \beta/\pi = \cos^{-1}\frac{1}{4}/\pi = .460106$.

(2). Let chord $OM = \frac{1}{2}a$, then in order that he may step out the third step he must step somewhere on the arc CM or its equal on the opposite side

$$\begin{aligned} \angle CAM &= \delta = \pi - (\beta + OAM) \\ &= \cos^{-1} \left(\frac{31\sqrt{105} - 7}{64} \right). \end{aligned}$$

$P_1 =$ chance he steps on this arc $= \delta/\pi = .379034$.

If his second step places him on arc CM then his third step must place him on the arc GKH . The $\angle KFH$ may vary from 0 to $\cos^{-1}(-\frac{1}{4})$.

$\therefore p_1 =$ chance that he steps on arc $GKH = \frac{\cos^{-1}(-\frac{1}{4})}{2\pi}$.

$\therefore p_1 = .304086$.

Now $p = P_1 \times p_1 = \delta/\pi \times \frac{\cos^{-1}(-\frac{1}{4})}{2\pi} = .115259$.

Solved with a different result by CHAS. C. CROSS.

55. Proposed by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics in Russell College, Lebanon, Va.

It has been clear for 15 consecutive days, what is the chance of the 16th day being cloudy?

Solution by the PROPOSER.

Let $p =$ chance, $p_1 =$ chance that 16th day is clear.

$$\therefore p_1 = \frac{\int_0^1 x^{1.6} dx}{\int_0^1 x^{1.5} dx} = \frac{1}{1.7}, \quad \therefore p = 1 - p_1 = \frac{1}{1.7}.$$

